

Four-dimensional Matter

A.M. Gevorkian , R. A. Gevorkian*

Institute of Physical Research, 378410, Ashtarak-2, Republic of Armenia

**Institute of Radiophysics and Electronics, 378410, Ashtarak-2, Republic of Armenia*

Abstract

Four-dimensional mass is determined in four-dimensional pseudo-Euclidean space as a physical invariant of that space. That invariant is discussed as an invariant of electromagnetic type. Finally, equations of Maxwell type are obtained for mass fields.

In Okun's publication "What is mass?" [1] the formula

$$m^2 = \frac{U^2}{c^4} - \frac{P^2}{c^2} \quad (1)$$

and comments on it looked very interesting to us for two reasons:

1) as a four-dimensional matter (in [1] it is defined as "simple mass").

The analysis of non-stationar phenomena and processes lead V.A. Ambartsumian to the idea of a possible new state of matter. In works [2], [3] and [4] the proto-star matter is discussed as a possible presentation of this state.

There are also a few works by R. Muradyan ([5] and [6]) on unusual manifestations (quantum) of macroscopic star systems (two-dimensional hadron).

Publication [1] notes that during fragmentation four-dimensional mass is not additive. The process of fragmentation continues to the point of occurrence of elementary particles (electron, proton) with emission of energy. Later the usual three-dimensional matter is constructed from those particles, which is the process of three-dimensionalization. The simplest form of this matter is the hydrogen atom. The further cooling of the environment by means of condensation, leads to the formation of usual star matter and its systems, and these systems may be unstable. In that case the famous disagreement between Ambartsumian and his opponents regarding the concept of formation of star systems becomes unimportant. We would like to remind that the first member of expression (1) is a square of a scalar, and the second member is a scalar - a module of square of a three-dimensional vector.

For compensation of the deficit of the mean cosmic density of matter some scientists proposed assumptions such as the "spontaneous birth of matter" (Hoyle), "existence of galaxies of neutrino matter" (Novikov), or existence of dark matter due to objects like the "black holes". Ambartsumian expressed opinions regarding this in [3], [4] and during seminars where one of the authors was present, that the law of energy conservation in its present form is incomplete. With respect to that we assume that the invariant (1) is a scalar of anti-symmetric tensor, which is created on the basis of three-dimensional (polar and axial) vectors.

2) as Lorentz invariant of square form with anti-symmetric tensor of four-dimensional matter [7].

$$M_{ik} = \begin{vmatrix} 0 & U_x & U_y & U_z \\ -U_x & 0 & -P_z & P_y \\ -U_y & P_z & 0 & -P_x \\ -U_z & -P_y & P_x & 0 \end{vmatrix} \quad (2)$$

Elements of this tensor can be considered as components of electric mass (polar three-dimensional vector) and magnetic mass (axial three-dimensional vector) fields.

With the help of this tensor we can construct invariant values with respect to Lorentz modifications. These invariants have the following form:

$$M_{ik}M^{ik} = I_1, \quad e^{iklm}M_{ik}M_{lm} = I_2 \text{ or } P^2 - U^2 = I_1, \quad UP = I_2. \quad (3)$$

I_1 corresponds to expression (1) and is discussed as Lagrangian of the physical system and satisfies Lagrange equation which transforms into a system of equations of Maxwell type. The four-dimensional form of these equations is:

$$\frac{\partial}{\partial x^i}M^{ik} = j^k, \quad \varepsilon^{iklm}\frac{\partial}{\partial x^k}M_{lm} = 0 \quad (4)$$

and in three-dimensional form it is

$$\begin{aligned} \operatorname{div} P &= 0 & \operatorname{rot} U &= \frac{1}{c} \frac{\partial P}{\partial t} \\ \operatorname{div} U &= \rho & \operatorname{rot} P &= \frac{1}{c} \frac{\partial U}{\partial t} + \frac{1}{c} j \end{aligned} \quad (5)$$

where ρ is the source (charge) of electric-mass static field which has the measurement of gr/cm and represents mass per unit length. And finally the wave equation in four-dimensional form is $\square A = 0$, where A is the vector-potential of the electromagnetic-mass field, and $\square = -\frac{\partial^2}{\partial x_i \partial x_k}$.

The source of field can be a mass string, as well as a mass ring. The movement of mass along a string or a ring (any locked configuration) induces magnetic-mass field of particular topology (toric, flat, etc).

It might be very interesting to place this in electric or magnetic environments, for instance inside an electric charge. A similar thing exists in the work of Ciama [6] in which he gives up the advantage of general covariance in favor of the vector theory of gravitation. The parameters of such an object can be considered when a quantum (calibration) theory is created similar to quantum electrodynamics or theory of electro-weak interactions. It is not excluded that these ideas may be virtual and may not have anything in common with real physics.

Literature

1. L. B. Okun "What is mass?" Physics - Uspekhi, 43(12) 2000.
2. Ambartsumyan V. A. Reports of the Academy of Science of Arm. SSR., 16, 97, 1953.
3. Ambartsumyan V. A. "Phenomena of Continuous Emission and Sources of Star Energy", Notices of the Byurakan Observatory, 13, 1954.
4. Ambartsumyan V. A. "Stars of T-Taurus and UV-Whale Type and the Phenomenon of continuous emission", Theses for the Symposium on Non-stationar Stars. Theses for the 9th Congress of the International Astronomic Society, 1955, 5, M., 1955.
5. Muradyan R., Astrophysics, 11, 237, 1975.
6. Muradyan R., Astrophysics, 13, 63, 1975.
7. Landau L. D., Lifshits E. M., "Field Theory", 1988.
8. Ciama D. W Roy Astron. Soc Monthly Notices 113, 34, 1953.

This figure "Gevorkian4.png" is available in "png" format from:

<http://arXiv.org/ps/physics/0106011v1>